BRL TR-3282

AD-A242 586



TECHNICAL REPORT BRL-TR-3282

BRL

A COMPARATIVE STUDY OF THE REACTION DYNAMICS OF A MODEL SYSTEM USING DIFFERENT CRITERIA IN PARAMETERIZING THE POTENTIAL ENERGY FUNCTION

> BETSY M. RICE CARY F. CHABALOWSKI GEORGE F. ADAMS RICHARD C. MOWREY MICHAEL J. PAGE

> > OCTOBER 1991

APPROVED FOR PUBLIC RELEASE; DISTRIBUTION IS UNLIMITED.

U.S. ARMY LABORATORY COMMAND

BALLISTIC RESEARCH LABORATORY
ABERDEEN PROVING GROUND, MARYLAND

91-15881

NOTICES

Destroy this report when it is no longer needed. DO NOT return it to the originator.

Additional copies of this report may be obtained from the National Technical Information Service, U.S. Department of Commerce, 5285 Port Royal Road, Springfield, VA 22161.

The findings of this report are not to be construed as an official Department of the Army position, unless so designated by other authorized documents.

The use of trade names or manufacturers' names in this report does not constitute indorsement of any commercial product.

UNCLASSIFIED

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average. I hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for information Operations and Reports, 1215 Jefferson Davis Highway Suite 1204 Attination 74, 22202-4302, and to the Office of Management and Budget Pagermyor Reduction Project (10740-1388) Washington DC 20503

Davis Highway, Suite 1204, Arlington, VA 22202-4302.	and to the Office of Management and	Budget, Paperwork Reduction Proj	ect (0704-0188), Washington, DC 20503
1 AGENCY USE ONLY (Leave blank)	2. REPORT DATE	3. REPORT TYPE AN	
4. TITLE AND SUBTITLE	October 1991	Final, Feb 91 - I	Mar 91 5. FUNDING NUMBERS
			5. FUNDING NUMBERS
A Comparative Study of the Reaction Dynamics of a Model System Using Different Criteria in Parameterizing the Potential Energy Function			PR: 1L161102AH43
6. AUTHOR(S)			
Betsy M. Rice, Cary F. Chab Richard C. Mowrey, and Micl		ns,	
7. PERFORMING ORGANIZATION NAME			8. PERFORMING ORGANIZATION
			REPORT NUMBER
		!	
		i	
9. SPONSORING / MONITORING AGENCY	NAME(S) AND ADDRESS(ES	3	10. SPONSORING / MONITORING
		•	AGENCY REPORT NUMBER
US Army Ballistic Research Laboratory			
ATTN: SLCBR-DD-T	•		BRL-TR-3282
Aberdeen Proving Ground, M	ID 21005-5066		
11. SUPPLEMENTARY NOTES			
Richard C. Mowrey and Mich	iael J. Page are employe	ees of the Naval Rese	earch Laboratory, Washington,
D.C. Currently in press for p	ublication in Chemical Pl	hysics Letters.	
12a. DISTRIBUTION / AVAILABILITY STAT	EMENT		12b. DISTRIBUTION CODE
Approved for public release;	distribution is unlimited.		
•			
13. ABSTRACT (Maximum 200 words)			
A comparative study of th	ie reaction dynamics of a	a model system using	a potential energy
function that has been param	neterized according to two	o different criteria is p	resented. The molecule
studied is methylene nitramin	e CH ₂ NNO ₂ and the read	ction studied is the N-	N bond scission reaction.
A mathematical description of parameters for the individual	terms in the model were	face for the system w	vas developed and
computations. The first mode	el was fitted to the Carte	illeu iu uaia ubiame sian second derivative	o ithe energy for
equilibrium methylene nitrami	ne, while the second mo	del was fitted using the	ne computed harmonic
frequencies for the molecule.	Unimolecular decay cur	rves for the N-N bond	scission reaction were
computed at six energies beto	ween 2.9384 and 4.1884	eV. Not only do the	two models disagree in
the computed reaction rates, but the rates predicted by Model 1 are first-order and time independent while those obtained with Model 2 indicate erratic and nonstatistical behavior. This study shows the			
While those obtained with mod	del 2 indicate erratic and	nonstatistical behavio	or. This study shows the
sensitivity of dynamics calcula different aspects of electronic	Ations to quantitative unit	erences in potential el	nergy surfaces fitted to
amorain appeals of stockers.	Structure Carculations.		
14. SUBJECT TERMS			15. NUMBER OF PAGES
			26
methylene nitramine; potentia	I energy surface; molecu	ılar dynamics;	16. PRICE CODE
unimolecular decomposition;	potential energy; energy		
	SECURITY CLASSIFICATION OF THIS PAGE	19. SECURITY CLASSIFIC OF ABSTRACT	CATION 20. LIMITATION OF ABSTRACT
	NCLASSIFIED	UNCLASSIFIED	UL

INTENTIONALLY LEFT BLANK.

TABLE OF CONTENTS

		<u>Page</u>
	LIST OF FIGURES	v
	LIST OF TABLES	v
	ACKNOWLEDGMENTS	vii
1.	INTRODUCTION	1
2.	POTENTIAL ENERGY SURFACE	2
2.1 2.2	Form of the Potential	_
3.	DISCUSSION	9
4.	REFERENCES	15
	DISTRIBUTION LIST	17

Ara ess :	on For	
311 G		K
Jacifi	Flu #d	Ö
Ву		
Distrib	bility all each	Codes
	Special	
11-1		

INTENTIONALLY LEFT BLANK.

LIST OF FIGURES

<u>Figure</u>		<u>Page</u>
1.	Structure of Methylene Nitramine at the Equilibrium Geometry Obtained at the 8-in-8 CASSCF/DZ Level	7
2.	Depictions of the Normal Modes of Vibrations of Methylene Nitramine for Modes (a) 4, (b) 11, and (c) 12 (See Table 3)	8
3.	Unimolecular Decay Curves for the N-N Bond Scission Reaction of Methylene Nitramine at (a) 2.9384 eV (b) 3.1884 eV, (c) 3.4384 eV, (d) 3.6884 eV, (e) 3.9384 eV, and (f) 4.1884 eV Total Energy	11
4.	Plot of the First-Order Decay Coefficient as a Function of Energy for (a) Model 1 and (b) Model 2 (See Text)	12
	LIST OF TABLES	
<u>Table</u>		Page
1.	Potential Energy Functions and Parameters	4
2.	Structural Parameters for CH ₂ NNO ₂ at the Equilibrium Geometry	5
3.	Normal Mode Frequencies of CH ₂ NNO ₂	6
4.	Computed Rate Coefficients	13

INTENTIONALLY LEFT BLANK.

ACKNOWLEDGEMENTS

The work at the Naval Research Laboratory (NRL) was supported by the Office of Naval Research through NRL; the ONR Physics Division (RCM), Contract N00014-91-WX-24193; and the ONR Mechanics Division (MP). BMR, CFC, and GFA acknowledge the assistance of the computer support staff of the U.S. Army Ballistic Research Laboratory (BRL) and equipment provided by the Productivity Enhancement Capital Investment Program.

INTENTIONALLY LEFT BLANK.

1. INTRODUCTION

The desire to employ classical mechanical molecular modeling and molecular dynamics for ever more complicated molecular systems has created the need for more general and accurate descriptions of the potential energy. Major goals of both molecular mechanics and molecular dynamics calculations are to predict and understand the behavior of the system at a molecular level. The success or failure in attaining these goals depends mainly on the accuracy of the potential energy function in describing the system. Often only limited data are available to use in developing these functions, such as vibrational frequencies and activation energies from experiment, or selected energies and geometries calculated by ab initio methods.

Molecular mechanics calculations employ force fields which are usually tailored to describe conformational changes that lie within a few kcal/mol of the equilibrium structure and are transferable among molecules. Recently developed force fields, such as MM3 (Allinger, Yuh, and Lii 1989) and CFF89 (Maple et al. 1990), provide more general descriptions of materials than did the first generation of force fields. While these force fields offer the opportunity to model well the properties of materials, there is an increased requirement for molecular information from which force field parameters can be derived. In several recent publications, force field parameters have been derived by fitting to computed properties and to the energy derivatives (first and second) calculated by ab initio quantum chemical techniques (Maple, Dinur, and Hagler 1988; Hagler et al. 1989; Dinur and Hagler 1989a, 1989b, 1990). The fitting techniques depend upon the application and upon the desire to derive transferable force field parameters. Inclusion of the additional information obtained from the ab initio calculations has been reasonably successful in deriving force fields that better describe the test molecule and related molecules.

In a molecular dynamics simulation of a molecular reaction, by contrast, the energy expression is generally designed for the specific problem, such as a decomposition path. In that case, the potential energy function is designed to describe the reactant and product, and, if applicable, the transition state for the reaction. Despite the difference in focus, the techniques that have proved successful in the development of force fields for molecular

models should have application in the derivation of a molecular dynamics potential energy function.

To test this assertion, we construct potential energy expressions for the methylene nitramine molecule, CH₂NNO₂, and examine the effects of these models on the dynamics of unimolecular decomposition via N-N bond rupture. Ab initio calculations for methylene nitramine, the transition state leading to the elimination of HONO and the N-N bond fragmentation path have been reported (Mowrey et. al 1990). In the present work, we assumed a specific molecular-mechanics-type functional form for the potential energy of CH₂NNO₂, and the parameters in this form were adjusted in two different ways. In the first, Model 1, the parameters were fit to the matrix of second derivatives of the energy with respect to Cartesian coordinates computed in Mowrey et al. (1990). The parameters for Model 2 were fitted only to the harmonic vibrational frequencies reported by Mowrey et al. (1990). For this study, the properties were computed for the equilibrium structure as determined using an 8-in-8 CASSCF with a double zeta atomic basis set.

In Section 2, we describe the functional form of the potential energy surface and comment on the features. The following section contains the description of the molecular dynamics calculations for both potential energy models and discusses the differences between the two sets of results.

2. POTENTIAL ENERGY SURFACE

2.1 Form of the Potential. The function used to describe methylene nitramine is :

$$V_{CH_{2}NNO_{2}} = \sum V_{CH}^{Stretch}(\tau_{i}) + V_{CN}^{Stretch} + V_{NN}^{Stretch} + \sum V_{NO}^{Stretch}(\tau_{i}) + \sum V_{HCH}^{Bend}(\tau_{i})$$

$$+ V_{ONO}^{BEND} + \sum V_{HCN}^{BEND}(\tau_{i}) + \left[\sum V_{NNO}^{BEND}(\tau_{i}) + V_{CNN}^{BEND} + \sum V_{CNNO}^{Torsion} + \sum V_{HCNN}^{Torsion} + V_{NCN}^{Wag} + V_{NNO_{2}}^{Wag}\right] F(R_{NN}) ,$$

$$(1)$$

where all of the interaction potential energy functions and parameters are given in Table 1 and $F(R_{NN})$ is a function which attenuates interaction terms as N-N dissociation occurs. The form of $F(R_{NN})$ is

$$F(R_{NN}) = \begin{bmatrix} 1.0, & R_{NN} \leq R_e^{NN} \\ \exp(-\alpha_{NN}[R_{NN} - R_e^{NN}]^4), & R_{NN} > R_e^{NN} \end{bmatrix}$$
(2)

where α_{NN} and R_e^{NN} are the Morse-function parameters for the N-N bond listed in Table 1. Some of the terms in Equation 1 show dependence on τ_i or $\dot{\tau}_i$ (the HCNN or CNNO dihedral angles, respectively). The force constants and geometries differ for these interactions depending on the orientation of these dihedral angles. The changes in the two types of parameters can be described as follows:

$$P = P^{cis} + (P^{trans} - P^{cis}) [1.0 - \cos(\tau)]/2$$
 (3)

where τ is the appropriate CNNO or HCNN dihedral angle and P denotes a force constant or geometric parameter. The terms V_{ijkl}^{Wag} are functions of the out-of-plane wagging angle, ω . The angle ω is given as

$$\cos(\omega) = \frac{(\overrightarrow{R_{ji}} \times \overrightarrow{R_{jk}})}{R_{ji}R_{ji}R_{jk}\sin(\theta_{kjl})} \cdot \overrightarrow{R_{ji}}$$
 (4)

and represents the angle between a bond ji and a vector perpendicular to the plane defined by two bonds jk and jl, with $\omega=90^\circ$ corresponding to the equilibrium configuration. The potential energy term for the wagging motion is the same as for the bending motion. Parameters for Model 1 were obtained by a non-linear least squares fit to the Cartesian second energy derivatives calculated by ab initio methods (Mowrey et al. 1990). Starting with the parameter set for Model 1, the parameters for Model 2 were adjusted by non-linear least squares to fit the ab initio frequencies only, without regard to the accompanying eigenvectors.

2.2 <u>Features of the Two Models</u>. The equilibrium geometries and normal mode frequencies for the two models are shown for comparison with the ab initio results in Tables 2 and 3, respectively. We have chosen the ab initio frequencies and eigenvectors as our standard for comparison. Because the parameters of the potential function for Model 2 were

Table 1. Potential Energy Functions and Parameters

			N _{Str}	$V^{Stretch} = D\{1 - \exp(-\alpha[R-R_a])\}^2 - D$.(-a[R-R _a])}² -	O			
Bond	9)Q	D(eV)	ື້ນ	R _a s(Å)	R ^{rans} (Å)	α^{ds}	α ^{ds} (Å-¹) ⁸	$\alpha^{\text{trans}}(A^{-1})^{\mathbf{a}}$	Å-¹)ª
						Model 1	Model 2	Model 1	Model 2
H S	3.9(3.9028	1.0	1.0734	1.0707	2.2477	2.2564	2.2739	2.2590
S S	5.2	989	7.	5209		2.5363	2.2408		
Z	1.5	1.5118	1.4	1141		2.4589	3.5673		
ON	4.0	4.0459	1.8	3034	1.2404	2.1330	2.2617	2.7129	1.7128
			V ^B	$V^{\text{Bend or Wag}} = K(\gamma \gamma_{\text{E}})^2$, $\gamma = \theta$ or ω	$\gamma_{\rm E}$) ² , $\gamma = \theta$ or	3			
Angle	k ^{ds} (eV-	k ^{ds} (eV-deg⁻²)ª	K ^{trans} (e)	k ^{rans} (eV-deg ⁻²) ^a	γ ^{ds} (deg)	Yrans	γ ^{rans} (deg)		
	Model 1	Model 2	Model 1	Model 2					
HCH	1.2834	0.6344			119.9159				
e HCN H	2.3974	2.1007	2.3738	3.5110	124.1249	115.	115.9592		
CNN	3.4250	5.7410			115.7203				
ONN	3.6293	15.4549	2.5295	9.4748	121.0758	114	114.4648		
° ONO	3.5308	2.7348			124.4595				
NNO ₂	1.2606	1.2606			90.000				
U C H	0.5351	0.5351			90.0000				
				$V^{\text{Torsion}} = k[1-\cos^2 t]$	[1-cos ² t]	,			
				Angle	k(eV) ^d				
				CNNO	0.0671				
				HCNN	0.9800				

^a Parameter allowed to vary in fitting.

Bond angle.
Cout-of-plane wag angle.
Same for both models.

Table 2. Structural Parameters for CH2NNO, at the Equilibrium Geometry

Internal ^a Coordinate	Ab initio⁵	Model 1	Model 2
R(CN) R(NN) R(NO) ^c R(NO) ^d R(CH) ^e R(CH) ^t	1.271 1.441 1.303 1.240 1.073 1.071	1.271 1.441 1.303 1.239 1.073 1.071	1.271 1.441 1.303 1.239 1.073 1.071
θ(CNN) θ(NNO) ^c θ(HCN) ^g θ(HCN) ^f θ(ONO)	115.700 121.100 114.500 124.100 116.000 124.400	115.700 121.100 114.500 124.100 116.000 124.500	115.700 121.100 114.500 124.100 116.000 124.500

Bond distances and angles are given in units of A and degrees, respectively.

determined by fitting to the ab initio frequencies, the agreement between the two is very close. This is the best fit to the frequencies that the functional form would allow. The comparison between the ab initio frequencies and those calculated from Model 1 is, as expected, less satisfactory. In fact, the frequencies calculated from Model 1 are in significant disagreement with the ab initio value for modes 2, 4, 6, and 11. The eigenvectors of Model 1, however, fairly well resemble those of the ab initio eigenvectors and are easily assignable for all modes. In contrast, there are 6 modes in Model 2 (6, 8, 10, 11, 12, and 13) for which the eigenvectors bear little resemblance to the corresponding standard. Of these, modes 8, 10, 11, and 12 cannot be assigned unambiguously to any of the ab initio eigenvectors. Mode 6, however, bears a slight resemblance to the ab initio eigenvector for mode 5. To illustrate, the eigenvectors corresponding to modes 4, 11, and 12 calculated by the 2 models are shown for comparison with the standards in Figure 2. Figure 2a shows the worst agreement in frequency between Model 1 and the standard. Inspection of this figure shows that the eigenvectors of both of the models are in relatively good agreement with the standard.

^b 8-in-8 CASSCF/DZ (Mowrey et. al 1990).

^c Oxygen atom no. 1 (see Figure 1).

^d Oxygen atom no. 2 (see Figure 1).

^e Hydrogen atom no. 1 (see Figure 1).

^f Hydrogen atom no. 2 (see Figure 1).

Table 3. Normal Mode Frequencies of CH₂NNO₂^a

Mode No.	Ab initio ^b	Model 1 ^c	Model 2 ^d
1	100	101	101
2	391	274	391
3	561	560	560
4	621	441	619
5	665	610	666
6	872	732	868
7	896	894	894
8	1220	1206	1220
9	1261	1263	1263
10	1311	1312	1311
11	1580	1447	1580
12	1612	1637	1613
13	1839	1839	1840
14	3361	3362	3356
15	3483	3492	3487
ZPE° (eV)	1.2259	1.1884	1.2255

^a Frequencies given in cm⁻¹.

Figure 2b again demonstrates a case where the frequency of the mode predicted by Model 1 disagrees with the standard, but the eigenvector is clearly assignable. In this case, however, the eigenvector of Model 2 does not resemble the standard in spite of its excellent agreement in frequency. In the last case, illustrated in Figure 2c, the frequencies from both models compare well with the standard, but the eigenvector of Model 2 is unassignable to this mode. The differences in vibrational frequencies between Model 1 and the standard merely reflect the inability to perfectly fit the matrix of second-derivatives with respect to energy for Model 1 which indicates deficiencies in the form of the potential energy function. On the other hand, the inability to reproduce the eigenvectors from Model 2 is due not only to deficiencies in the functional form, but also to the fact that the parameters for Model 2 were fit to less potential energy surface information, specifically concerning the shape of the surface.

^b 8-in-8 CASSCF/DZ (Mowrey et al. 1990).

^c Eigenvectors corresponding to these frequencies matched to ab initio eigenvectors.

Frequencies matched to ab initio frequencies regardless of eigenvector form.

^e Zero point vibrational energy.





Figure 1. <u>Structure of Methylene Nitramine at the Equilibrium Geometry Obtained at the 8-in-8 CASSCF/DZ Level.</u>



Ab initio results are shown in the first column of the figures, while the results for Models 1 and 2 are shown in the second and third columns, respectively. The harmonic frequencies of the modes are listed in parentheses.

Figure 2. Depictions of the Normal Modes of Vibrations of Methylene Nitramine for Modes
(a) 4, (b) 11, and (c) 12 (See Table 3).

In almost all instances, there will be deficiencies in the form of the potential energy function; therefore, it will be difficult to attain good fits. In addition to defining a suitably flexible function, a modeler is also faced with the issue of whether it is better to correctly describe the eigenvectors at the sacrifice of agreement in frequencies or to fit to the frequencies only, thereby ignoring the detailed motion of the atoms. For this system, the assumption that Model 2 accurately describes the system under study based only on agreement with the harmonic frequencies is deceptive, considering how poorly some of the eigenvectors compare with the ab initio values. Although differences in the dynamics occur near the equilibrium geometry due to the differences in the fitting criteria for the model (i.e., the normal modes), we wish to further explore how the reaction dynamics differ depending on the different criteria used in fitting the functions. To investigate this, we performed a series of molecular dynamics calculations on the two models; specifically, we investigated the unimolecular decomposition of CH₂NNO₂.

3. DISCUSSION

Although there are experimental evidence (Zhao, Hintsa, and Lee 1988) and theoretical predictions (Mowrey et. al 1990) that unimolecular decay of CH₂NNO₂ is more likely to occur by concerted dissociation pathways than through N-N bond scission, for our purposes our models only describe bond scission reactions. The N-N bond is the weakest bond in this system by 2.4 eV, and this was the only reaction we observed.

Ensembles of trajectories at six energies were integrated using a variable step size Adams-Moulton fourth-order predictor corrector integrator (Miller and George 1972). Relative error tolerance was set at 10⁻⁷. Before each ensemble of trajectories is integrated, the system (including the zero point energy) is equipartitioned among the normal modes in the form of kinetic energy. A warm-up trajectory of approximately 0.03 ps is performed, and a Markov walk of 75,000 steps is taken to randomize the energy of the system. A trajectory is then integrated until the N-N bond exceeds 6.0 Å or until the trajectory integration exceeds a maximum time of 30 ps. A sequence of 1,500 Markov moves is taken from the starting point of the previous trajectory and the integration/Markov walk pattern is repeated until 1,000 trajectories have been integrated. According to the standard Metropolis Monte Carlo sampling procedure (Metropolis et. al 1953; Raff and Thompson 1985; Brady, Doll, and


Thompson 1981), if a trial move taken from the starting point of a previous trajectory j is rejected, then the result of the previous trajectory is counted again and recorded as the j+1 contribution to the ensemble average. The present Metropolis procedure was designed such that approximately 50% of all attempted moves were rejected. Because of this rejection ratio, results from approximately 50% of the 1,000 trajectories integrated were recounted and included in the decay curves. In the Markov walk, the system is not allowed to exceed an N-N bond distance of 3.75 Å. Each trajectory included in the decay curves was required to undergo at least one N-N bond vibration before dissociation; if it did not, it was not included. The time of decomposition was marked at the last inner turning point of the N-N bond vibration.

Figure 3 shows unimolecular decay curves of ln(P) vs. time where P is the fraction of undissociated trajectories at time *t* for Models 1 and 2 at 2.9384, 3.1884, 3.4384, 3.6884, 3.9384, and 4.1884 eV. First-order rate coefficients were extracted from the straight line portions of the curves and are listed in Table 4. In all of the figures, the decay curves of Model 1 are well fit to a straight line, indicating the rate in this system is first-order and time independent. Also, as expected, the rate coefficients show a monotonic increase with increasing energy. Another indication of the well-behaved nature of this model is that the rates can be described by the RRK statistical model (Robinson and Holbrook 1972)

$$k(E) = A \left[\frac{E - E_o}{E} \right]^{s-1}, \tag{4}$$


where $A = 5.1 \text{ ps}^{-1}$, $E_o = 2.7 \text{ eV}$ and s = 1.3. The calculated rates from Model 1 and those predicted from Equation 4 are shown in Figure 4a.

The decay curves for Model 2 are not well behaved; in Figures 2b, 2d, and 2e there are at least two regions that are linear but differ in slope. The slopes for the initial regions of these three curves are significantly steeper than for the region of longer lifetimes, indicating that there is a faster reaction rate for the early decomposition events and a slower rate for longer reaction times. Like Model 1, the three other decay curves (Figures 2a, 2c, and 2f) can be reasonably fit to a straight line. We attempted to fit the data from Model 2 to Equation 4 in



Results from Model 1 are denoted by filled circles and results from Model 2 are denoted by crosses.

Figure 3. Unimolecular Decay Curves for the N-N Bond Scission Reaction of Methylene Nitramine at (a) 2.9384, (b) 3.1884 eV, (c) 3.4384 eV, (d) 3.6884, (e) 3.9384 eV, and (f) 4.1884 eV Total Energy.



The filled circles are the rates extracted from the trajectory results and the curves are the results of the best fit of these results to the RRK expression, Equation 4.

Figure 4. Plot of the First-Order Decay Coefficient as a Function Energy for (a) Model 1 and (b) Model 2 (See Text).

Table 4. Computed Rate Coefficients^a

Energy (eV) ^b	k (ps ⁻¹)		
	Model 1	Mod	el 2°
		Fast	Slow
2.94	1.64	3.96	
3.19	2.54	1.46	0.142
3.44	2.98	4.36	
3.69	3.23	3.71	1.310
3.94	3.48	4.80	0.729
4.19	3.57	5.38	

The rate coefficients k are extracted from the least squares fits of the decay curves to the model of first order reaction, ln(p) = -kt (Figure 3).

the same manner as for Model 1 using the rate coefficients extracted from the curves in Figures 2a, 2c, and 2f and the "fast" region of the curves in Figures 2b, 2d, and 2e. Because Figures 2a, 2c, and 2f did not unambiguously exhibit a "slow" region of the curves, we did not attempt to fit the "slow" rates to Equation 4. Comparison of the best fit of Equation 4 to the rates extracted from Model 2 are shown in Figure 4b; the best fit parameters for this equation are 11.9 ps⁻¹, 1.178 eV and 2.97 for A, E_o , and s, respectively. As shown strikingly in Figure 4b and Table 4, the rates from Model 2 have an erratic energy dependence, and are poorly described by the RRK theory. These, coupled with bimodal behavior of the decay curves in Figures 2b, 2d, and 2e, are indicative of nonstatistical behavior, possibly due to bottlenecks in the system (Hase 1976).

Because CH₂NNO₂ has never been isolated and unimolecular decay experiments have never been done on this molecule, particularly this channel, we cannot state unequivocally which model more accurately represents the true system. We have shown, however, that with little effort, nonstatistical behavior can be incorporated into a model, and at the very least, we have shown that the rates of decomposition differ between the two models at all energies studied. We wish to stress that Model 2, which predicts nonstatistical behavior, is fit to the observable, the normal mode frequencies. If this nonstatistical behavior is only an artifact of the fitting procedure, one could easily be led to erroneous conclusions about a given system

Includes the zero point energy.

^c See text.

which exhibits very interesting dynamical behavior. The erratic nature of the dynamics results obtained from Model 2 leads one to question the usefulness for prediction and understanding the system as described by Model 2.

4. REFERENCES

- Allinger, N. L., Y. H. Yuh, and J. H. Lii. <u>Journal of the American Chemical Society</u>, vol. 111, p. 8551, 1989.
- Brady, J. W., J. D. Doll, and D. L. Thompson. <u>Journal of Chemical Physics</u>, vol. 74, p. 1026, 1981.
- Dinur, U., and A. T. Hagler. <u>Journal of the American Chemical Society</u>, vol. 111, p. 5149, 1989a
- Dinur, U., and A. T. Hagler. Journal of Chemical Physics, vol. 91, pp. 2949, 2959, 1989b.
- Dinur, U., and A. T. Hagler. Journal of Computational Chemistry, vol. 11, p. 1234, 1990.
- Hagler, A. T., J. R. Maple, T. S. Thatcher, G. B. Fitzgerald, and U. Dinur. <u>Computer Simulation of Biomolecular Systems-Theoretical and Experimental Applications</u>. Edited by W. F. van Gunsteren and P. K. Weiner. ESCOM, Leiden, pp. 149-167, 1989.
- Hase, W. L. "Dynamics of Molecular Collisions, Part B." <u>Modern Theoretical Chemistry</u>, vol. 2. Edited by W. H. Miller, New York: Plennum Press, pp. 121-169, 1976.
- Maple, J. R., U. Dinur, and A. T. Hagler. <u>Proceedings of the National Academy of Science USA</u>, vol. 85, p. 5350, 1988.
- Maple, J. R., T. S. Thatcher, U. Dinur, and A. T. Hagler. <u>Chemical Design Automation News</u>, vol. 5, no. 9, p. 10, 1990.
- Metropolis, N., A. W. Rosenbluth, M. N. Rosenbluth, A. H. Teller, and E. Teller. <u>Journal of Chemical Physics</u>, vol. 21, p. 1087, 1953.
- Miller, W. H., and T. F. George. <u>Journal of Chemical Physics</u>, vol. 56, p. 5668, 1972.
- Mowrey, R. C., M. Page, G. F. Adams, and B. H. Lengsfield III. <u>Journal of Chemical Physics</u>, vol. 93, p. 1857, 1990.
- Raff, L. M., and D. L. Thompson. <u>Theory of Chemical Reaction Dynamics</u>, vol. 1. Edited by M. Baer. Boca Raton, Florida: Chemical Rubber, pp. 1-121, 1985.
- Robinson, P. J., and K. A. Holbrook. <u>Unimolecular Reactions</u>. New York: John Wiley and Sons, Ltd., 1972.
- Zhao X., E. J., Hintsa, and Y. T. Lee. <u>Journal of Chemical Physics</u>, vol. 88, p. 801, 1988.

INTENTIONALLY LEFT BLANK.

No. of Copies	Organization	No. of Copies	Organization
2	Administrator Defense Technical Info Center ATTN: DTIC-DDA Cameron Station Alexandria, VA 22304-6145	1	Commander U.S. Army Missile Command ATTN: AMSMI-RD-CS-R (DOC) Redstone Arsenal, AL 35898-5010
1	Commander U.S. Army Materiel Command ATTN: AMCDRA-ST 5001 Eisenhower Avenue Alexandria, VA 22333-0001	1	Commander U.S. Army Tank-Automotive Command ATTN: ASQNC-TAC-DIT (Technical Information Center) Warren, MI 48397-5000
		1	Director
	Commander U.S. Army Laboratory Command ATTN: AMSLC-DL 2800 Powder Mill Road Adelphi, MD 20783-1145	1	U.S. Army TRADOC Analysis Command ATTN: ATRC-WSR White Sands Missile Range, NM 88002-5502 Commandant
	Commander U.S. Army Armament Research, Development, and Engineering Center		U.S. Army Field Artillery School ATTN: ATSF-CSI FI. SIII, OK 73503-5000
	ATTN: SMCAR-IMI-I Picatinny Arsenal, NJ 07806-5000	(Class. only)1	Commandant U.S. Army Infantry School ATTN: ATSH-CD (Security Mgr.)
	Commander U.S. Army Armament Research, Development, and Engineering Center	(Unclass. only)1	Fort Benning, GA 31905-5660 Commandant
	ATTN: SMCAR-TDC Picatinny Arsenal, NJ 07806-5000		U.S. Army Infantry School ATTN: ATSH-CD-CSO-OR Fort Benning, GA 31905-5660
	Director Benet Weapons Laboratory U.S. Army Armament Research,		Air Force Armament Laboratory ATTN: WL/MNOI
	Development, and Engineering Center ATTN: SMCAR-CCB-TL Watervliet, NY 12189-4050		Eglin AFB, FL 32542-5000 Aberdeen Proving Ground
(Unclass. only)1	Commander	2	Dir, USAMSAA
	U.S. Army Armament, Munitions and Chemical Command ATTN: AMSMC-IMF-L		ATTN: AMXSY-D AMXSY-MP, H. Cohen
	Rock Island, IL 61299-5000		Cdr, USATECOM ATTN: AMSTE-TC
	Director U.S. Army Aviation Research and Technology Activity ATTN: SAVRT-R (Library) M/S 219-3 Ames Research Center		Cdr, CRDEC, AMCCOM ATTN: SMCCR-RSP-A SMCCR-MU SMCCR-MSI
	Moffett Field, CA 94035-1000	1	Dir, VLAMO ATTN: AMSLC-VL-D
			Dir, BRL ATTN: SLCBR-DD-T

No. of	
<u>Copies</u>	Organization

- 1 HQDA (SARD-TR, C.H. Church) WASH DC 20310-0103
- 4 Commander
 US Army Research Office

ATTN: R. Ghirardelli

D. Mann R. Singleton

R. Shaw

P.O. Box 12211

Research Triangle Park, NC

27709-2211

2 Commander

US Army Armament Research,
Development, and Engineering Center
ATTN: SMCAR-AEE-B, D.S. Downs

SMCAR-AEE, J.A. Lannon

Picatinny Arsenal, NJ 07806-5000

1 Commander

US Army Armament Research,
Development, and Engineering Center
ATTN: SMCAR-AEE-BR, L. Harris

- Picatinny Arsenal, NJ 07806-5000
- 2 Commander US Army Missile Command ATTN: AMSMI-RD-PR-E, A.R. Maykut AMSMI-RD-PR-P, R. Betts

Redstone Arsenal. AL 35898-5249

- Office of Naval Research
 Department of the Navy
 ATTN: R.S. Miller, Code 432
 800 N. Quincy Street
 Arlington, VA 22217
- Commander
 Naval Air Systems Command
 ATTN: J. Ramnarace, AIR-54111C
 Washington, DC 20360
- Commander
 Naval Surface Warfare Center
 ATTN: J.L. East, Jr., G-23
 Dahlgren, VA 22448-5000

No. of Copies Organization

Commander
 Naval Surface Warfare Center
 ATTN: R. Bernecker, R-13
 G.B. Wilmot, R-16

Silver Spring, MD 20903-5000

5 Commander

Naval Research Laboratory

ATTN: M.C. Lin

J. McDonald

E. Oran

J. Shnur

R.J. Doyle, Code 6110

Washington, DC 20375

- Commanding Officer
 Naval Underwater Systems
 Center Weapons Dept.
 ATTN: R.S. Lazar/Code 36301
 - ATTN: R.S. Lazar/Code 3630 Newport, RI 02840
- Commander
 Naval Weapons Center
 ATTN: T. Boggs, Code 388
 T. Parr, Code 3895
 China Lake, CA 93555-6001
- Superintendent
 Naval Postgraduate School
 Dept. of Aeronautics
 ATTN: D.W. Netzer
 Monterey, CA 93940
- 3 AL/LSCF
 ATTN: R. Corley
 R. Geisler
 J. Levine
 Edwards AFB, CA 93523-5000
- 1 AL/MKPB ATTN: B. Goshgarian Edwards AFB, CA 93523-5000
- 1 AFOSR ATTN: J.M. Tishkoff Bolling Air Force Base Washington, DC 20332
- 1 OSD/SDIO/IST ATTN: L. Caveny Pentagon Washington, DC 20301-7100

1 Commandant USAFAS ATTN: ATSF-TSM-CN Fort Sill, OK 73503-5600

1 F.J. Seiler ATTN: S.A. Shackleford USAF Academy, CO 80840-6528

 University of Dayton Research Institute ATTN: D. Campbell AL/PAP Edwards AFB, CA 93523

NASA

 Langley Research Center
 Langley Station
 ATTN: G.B. Northam/MS 168
 Hampton, VA 23365

4 National Bureau of Standards ATTN: J. Hastie M. Jacox T. Kashiwagi H. Semerjian

US Department of Commerce Washington, DC 20234

 Aerojet Solid Propulsion Co. ATTN: P. Micheli Sacram. nto, GA 95813

 Applied Combustion Technology, Inc. ATTN: A.M. Varney
 P.O. Box 607885
 Orlando, FL 32860

Applied Mechanics Reviews
 The American Society of
 Mechanical Engineers
 ATTN: R.E. White
 A.B. Wenzel

 345 E. 47th Street
 New York, NY 10017

Atlantic Research Corp.
 ATTN: M.K. King
 5390 Cherokee Avenue
 Alexandria, VA 22314

No. of Copies Organization

Atlantic Research Corp.
 ATTN: R.H.W. Waesche
 7511 Wellington Road
 Gainesville, VA 22065

1 AVCO Everett Research
Laboratory Division
ATTN: D. Stickler
2385 Revere Beach Parkway
Everett, MA 02149

1 Battelle ATTN: TACTEC Library, J. Huggins 505 King Avenue Columbus, OH 43201-2693

Cohen Professional Services
 ATTN: N.S. Cohen
 141 Channing Street
 Redlands, CA 92373

 Exxon Research & Eng. Co. ATTN: A. Dean Route 22E Annandale, NJ 08801

 Ford Aerospace and Communications Corp.
 DIVAD Division
 Div. Hq., Irvine
 ATTN: D. Williams
 Main Street & Ford Road
 Newport Beach, CA 92663

1 General Applied ScienceLaboratories, Inc.77 Raynor AvenueRonkonkama, NY 11779-6649

 General Electric Ordnance Systems
 ATTN: J. Mandzy
 100 Plastics Avenue
 Pittsfield, MA 01203

 General Motors Rsch Labs Physical Chemistry Department ATTN: T. Sloane Warren, MI 48090-9055

- Hercules, Inc.
 Allegheny Ballistics Lab.
 ATTN: W.B. Walkup
 E.A. Yount
 P.O. Box 210
 Rocket Center, WV 26726
- 1 Alliant Techsystems, Inc. Marine Systems Group ATTN: D.E. Broden/ MS MN50-2000 600 2nd Street NE Hopkins, MN 55343
- Alliant Techsystems, Inc. ATTN: R.E. Tompkins MN38-3300 5700 Smetana Drive Minnetonka, MN 55343
- 1 IBM Corporation ATTN: A.C. Tam Research Division 5600 Cottle Road San Jose, CA 95193
- IIT Research Institute ATTN: R.F. Remaly
 West 35th Street Chicago, IL 60616
- Director

 Lawrence Livermore
 National Laboratory

 ATTN: C. Westbrook

 M. Costantino
 P.O. Box 808
 Livermore, CA 94550
- Lockheed Missiles & Space Co. ATTN: George Lo
 3251 Hanover Street
 Dept. 52-35/B204/2
 Palo Alto, CA 94304
- Director
 Los Alamos National Lab
 ATTN: B. Nichols, T7, MS-B284
 P.O. Box 1663
 Los Alamos, NM 87545

- National Science Foundation ATTN: A.B. Harvey Washington, DC 20550
- 1 Olin Ordnance ATTN: V. McDonald, Library P.O. Box 222 St. Marks, FL 32355-0222
- Paul Gough Associates, Inc. ATTN: P.S. Gough 1048 South Street Portsmouth, NH 03801-5423
- Princeton Combustion Research Laboratories, Inc. ATTN: M. Summerfield N.A. Messina 475 US Highway One Monmouth Junction, NJ 08852
- Hughes Aircraft Company ATTN: T.E. Ward 8433 Fallbrook Avenue Canoga Park, CA 91303
- Rockwell International Corp. Rocketdyne Division
 ATTN: J.E. Flanagan/HB02
 6633 Canoga Avenue
 Canoga Park, CA 91304
- Director

 Sandia National Laboratories
 Division 8354

 ATTN: R. Cattolica
 S. Johnston
 P. Mattern
 D. Stephenson
 Livermore, CA 94550
- Science Applications, Inc.
 ATTN: R.B. Edelman
 23146 Cumorah Crest
 Woodland Hills, CA 91364
- 3 SRI International
 ATTN: G. Smith
 D. Crosley
 D. Golden
 333 Ravenswood Avenue
 Menlo Park, CA 94025

- Stevens Institute of Tech. Davidson Laboratory ATTN: R. McAlevy, III Hoboken, NJ 07030
- Sverdrup Technology, Inc. LERC Group
 ATTN: R.J. Locke, MS SVR-2 2001 Aerospace Parkway
 Brook Park, OH 44142
- Sverdrup Technology, Inc. ATTN: J. Deur
 2001 Aerospace Parkway
 Brook Park, OH 44142
- Thiokol Corporation Elkton Division
 ATTN: S.F. Palopoli P.O. Box 241
 Elkton, MD 21921
- Thiokol Corporation
 Wasatch Division
 ATTN: S.J. Bennett
 P.O. Box 524
 Brigham City, UT 84302
- United Technologies Research Center ATTN: A.C. Eckbreth East Hartford, CT 06108
- United Technologies Corp.
 Chemical Systems Division
 ATTN: R.S. Brown
 T.D. Myers (2 copies)
 P.O. Box 49028
 San Jose, CA 95161-9028
- Universal Propulsion Company ATTN: H.J. McSpadden Black Canyon Stage 1 Box 1140 Phoenix, AZ 85029
- Veritay Technology, Inc.
 ATTN: E.B. Fisher
 4845 Millersport Highway
 P.O. Box 305
 East Amherst, NY 14051-0305

- Brigham Young University
 Dept. of Chemical Engineering
 ATTN: M.W. Beckstead
 Provo, UT 84058
- 1 California Institute of Tech. Jet Propulsion Laboratory ATTN: L. Strand/MS 512/102 4800 Oak Grove Drive Pasadena, CA 91109
- 1 California Institute of Technology ATTN: F.E.C. Culick/ MC 301-46 204 Karman Lab. Pasadena, CA 91125
- University of California
 Los Alamos Scientific Lab.
 P.O. Box 1663, Mail Stop B216
 Los Alamos, NM 87545
- University of California, Berkeley
 Chemistry Deparment
 ATTN: C. Bradley Moore
 211 Lewis Hall
 Berkeley, CA 94720
- University of California,
 San Diego
 ATTN: F.A. Williams
 AMES, B010
 La Jolla, CA 92093
- 2 University of California, Santa Barbara Quantum Institute ATTN: K. Schofield M. Steinberg Santa Barbara, CA 93106
- University of Colorado at Boulder
 Engineering Center
 ATTN: J. Daily
 Campus Box 427
 Boulder, CO 80309-0427

- University of Southern

 California
 Dept. of Chemistry

 ATTN: S. Benson

 Wittig

 Los Angeles, CA 90007
- Cornell University
 Department of Chemistry
 ATTN: T.A. Cool
 Baker Laboratory
 Ithaca, NY 14853
- University of Delaware ATTN: T. Brill Chemistry Department Newark, DE 19711
- University of Florida Dept. of Chemistry ATTN: J. Winefordner Gainesville, FL 32611
- 3 Georgia Institute of Technology School of Aerospace Engineering ATTN: E. Price W.C. Strahle B.T. Zinn Atlanta, GA 30332
- University of Illinois
 Dept. of Mech. Eng.
 ATTN: H. Krier
 144MEB, 1206 W. Green St.
 Urbana, IL 61801
- Johns Hopkins University/APL Chemical Propulsion Information Agency ATTN: T.W. Christian Johns Hopkins Road Laurel, MD 20707
- University of Michigan
 Gas Dynamics Lab
 Aerospace Engineering Bldg.
 ATTN: G.M. Faeth
 Ann Arbor, MI 48109-2140

- University of Minnesota Dept. of Mechanical Engineering ATTN: E. Fletcher Minneapolis, MN 55455
- 3 Pennsylvania State University Applied Research Laboratory ATTN: K.K. Kuo H. Palmer M. Micci University Park, PA 16802
- Pennsylvania State University
 Dept. of Mechanical Engineering
 ATTN: V. Yang
 University Park, PA 16802
- 1 Polytechnic Institute of NY Graduate Center ATTN: S. Lederman Route 110 Farmingdale, NY 11735
- Princeton University
 Forrestal Campus Library
 ATTN: K. Brezinsky
 I. Glassman
 P.O. Box 710
 Princeton, NJ 08540
- Purdue University
 School of Aeronautics
 and Astronautics
 ATTN: J.R. Osborn
 Grissom Hall
 West Lafayette, IN 47906
- 1 Purdue University Department of Chemistry ATTN: E. Grant West Lafayette, IN 47906
- 2 Purdue University School of Mechanical Engineering ATTN: N.M. Laurendeau S.N.B. Murthy TSPC Chaffee Hall West Lafayette, IN 47906

- Rensselaer Polytechnic Inst.
 Dept. of Chemical Engineering ATTN: A. Fontijn Troy, NY 12181
- 1 Stanford University
 Dept. of Mechanical
 Engineering
 ATTN: R. Hanson
 Stanford, CA 94305
- 1 University of Texas
 Dept. of Chemistry
 ATTN: W. Gardiner
 Austin, TX 78712
- University of Utah
 Dept. of Chemical Engineering
 ATTN: G. Flandro
 Salt Lake City, UT 84112
- Virginia Polytechnic Institute and State University ATTN: J.A. Schetz Blacksburg, VA 24061
- Freedman Associates
 ATTN: E. Freedman
 2411 Diana Road
 Baltimore, MD 21209-1525

INTENTIONALLY LEFT BLANK.

USER EVALUATION SHEET/CHANGE OF ADDRESS

This laboratory undertakes a continuing effort to improve the quality of the reports it

publishes. Your comments/answers below will aid us in our efforts. 1. Does this report satisfy a need? (Comment on purpose, related project, or other area of interest for which the report will be used.) 2. How, specifically, is the report being used? (Information source, design data, procedure, source of ideas, etc.) 3. Has the information in this report led to any quantitative savings as far as man-hours or dollars saved, operating costs avoided, or efficiencies achieved, etc? If so, please elaborate. 4. General Comments. What do you think should be changed to improve future reports? (Indicate changes to organization, technical content, format, etc.) BRL Report Number BRL-TR-3282 Division Symbol _____ Check here if desire to be removed from distribution list. Check here for address change. Current address: Organization Address **DEPARTMENT OF THE ARMY** Director NO POSTAGE U.S. Army Ballistic Research Laboratory ATTN: SLCBR-DD-T NECESSARY IF MAILED Aberdeen Proving Ground, MD 21005-5066 IN THE UNITED STATES **BUSINESS REPLY MAIL** OFFICIAL BUSINESS FIRST CLASS PERMIT No 0001, APG, MD Postage will be paid by addressee Director U.S. Army Ballistic Research Laboratory ATTN: SLCBR-DD-T

Aberdeen Proving Ground, MD 21005-5066